



Soybean Meal Information Center

FACT SHEET

Soybean Use - Aquaculture

PAUL B. BROWN
DEPARTMENT OF FORESTRY AND
NATURAL RESOURCES
PURDUE UNIVERSITY
WEST LAFAYETTE, IN 47907-1159

KEITH SMITH
KEITH SMITH AND ASSOCIATES
15 WINCHESTER ROAD
FARMINGTON, MO 63640



Soybean Meal in Aquaculture

There are lots of fish in the sea, as the saying goes. In fact, we know of more than 30,000 species.

But it will take more than the naturally occurring fish in the sea and commercial harvest alone to meet the future demand for fish as a foodstuff. In the late 1980s, commercial harvest was declared at maximum sustainable yield, prompting increased attention to “fish farming.”

Since 1990, most of the increases in aquaculture production have been with fish, mollusks and crustaceans in the southern hemisphere, primarily Asia and developing countries, with production nearly equally divided between freshwater and saltwater species.

Demand for new food aquaculture production during the next 30 to 40 years is expected to be between 60 and 120 million metric tons (MMT), rivaling the global poultry and swine industries. What will be the source of nutrition for these cultured fish?

Currently, 358 species of fish are cultured, of which 54 have been fed soybeans — the focus of nutritional research for new cultured species. But additional research is needed to confirm soy’s role in supplying protein to this new and growing market.



Species of Aquatic Animal

in Which Soybean Meal has Been Evaluated

Species

Number of Technical Reports in Literature

Atlantic salmon	13
Rainbow trout	41
Coho salmon	4
Chinook salmon	1
Tilapia	14
Ayu	1
Pacu	1
Siberian sturgeon	1
Largemouth bass	1
Hybrid striped bass	9
Striped bass	2
Turbot	1
Red drum	5
Tin foil barb	1
Giant gouramy	1
Yellowtail	7
Korean rockfish	2
Seabass	6
Seabream	10
Japanese sea bass	1
Milkfish	1
Bluntnose black bream	1
Australian snapper	2
Plaice	1
Winter flounder	1
Japanese flounder	3
Greenback flounder	1
African sole	1
Florida pompano	1
Channel catfish	13
African catfish	10
Blue catfish	1
Chinese longsnout catfish	1
Yellow mystus — tropical catfish	1
Indian carp — Labeo	5
Indian carp — Catla	3
Indian carp — Mrigal	2
Black carp	1
Grass carp	2
Common carp	16
Chinese hairy crab	1
Chinese mitten-handed crab	1
Pacific white shrimp	4
Grass shrimp	1
Banana shrimp	1
Indian white shrimp	2
Australian crayfish	4
Red swamp crayfish	4
Spiny lobster	1
American lobster	1
Green sea urchin	1
Abalone	2



Replacing Fish Meal and Fish Oil

Initial diets for aquaculture species typically contain high levels of fish meal and fish oil, which are flavorful ingredients for aquatic animals. Fish meal is a high protein ingredient with a good balance of quality essential amino acids (EAA), and fish oil contains n-3 fatty acids, required by many aquatic animals. Most aquatic animals grow well when fed relatively high levels of crude protein and lipid, and essential amino acid and fatty acid concentrations are priority considerations when formulating fish diets.

However, supplies of fish meal and oil are insufficient to realize needed growth in aquaculture production during the next 40 years. Global fish meal supplies are approximately 6.8 MMT annually, of which approximately 1.7 MMT, or 25 percent, is used in diets for aquatic animals. If we assume aquaculture grows the least amount predicted (60 million metric tons), and we can restrict fish meal usage to 10 percent of the diet, then aquaculture will demand 6 MMT of fish meal in the next 40 years. Keep in mind, the current global supply is only 6.8 MMT, and this will most likely not increase.

Fish meal contains higher crude protein concentrations than solvent-extracted soybean meal, but soybean meal prices are more stable than fish meal prices due to the more consistent supply of soy. Availability will likely be the primary deterrent to the use of fish meal in the future. Supplies and price are affected by such natural occurrences as El Niño, which will likely continue to bring uncertainty to future supplies of fish meal, as it did during the most recent cycle.

Aquaculture currently demands 0.4 MMT, or approximately 36 percent, of the annual global fish oil supply. As with fish meal, this is a finite resource, and increases in demand will strip supply if growth projections for aquaculture are realized. Replacing fish oil in aquatic diets will most likely be a more difficult challenge than replacing fish meal.

Dietary development for aquaculture has, in general, followed a consistent pattern. The first evaluations of new species, often unpublished, use those diets that are available. Most initial diet choices have been high fish-meal diets formulated for trout or salmon @ diets that are palatable by most species of fish. As studies progress, diets are gradually modified to more closely meet the nutritional requirements of target species.

Initial studies with a new species include evaluation of optimal dietary crude protein concentration, optimal protein to non-protein energy ratio, optimal ratio of carbohydrate to lipid and, finally, evaluation of commercial feedstuffs.

Is Soy the Answer?

In earlier nutrition work, as dietary development for ruminants and other animals progressed, essential amino acid requirements were recognized as being important.


However, of those aquatic species currently cultured, researchers have estimates of all 10 identified EAA requirements for only nine species. Lysine and methionine requirements, the two most limiting essential amino acids in feed formulations for most species, are known for another eight to 10 species. Despite this lack of specific knowledge, various forms of soybean meal have been evaluated in a variety of aquatic animals.



The table below shows species in which any form of soybean meal has been used as a source of crude protein and EAA in a diet. Many of these published reports are not formal evaluations of soy for target species, but rather studies of an initial diet that contains soy, or digestibility values using a form of soybean as a test feedstuff. The full-fat soybean meal designation includes raw, heated and roasted soybeans.

Slowly but surely, progress is being made. An article published in 2000 identified 17 species in which formal evaluations of

soybeans had been conducted. That number has now grown to 54 species. However, despite the number of evaluations, the specificity in the research is generally lacking. For example, there have been few definitive studies on trypsin inhibitors from soybeans and their effects on fish.





The article summarized the available data on the use of soy in fish, and argued that virtually all fish could handle a minimum of 10 percent to 15 percent soybean meal in their diets, but that several of the more carnivorous species could not handle more than 20 percent. The salmonids (trout, salmon and char) are one of the most sensitive species to soy, handling no more

than 25 percent to 30 percent, with some species handling no more than 15 percent soy. It is unclear why this limitation occurs.

With some of the newer aquaculture species, soybean incorporation into the diet can be relatively high. For example, once the critical EAA were established for the hybrid striped bass, researchers were able to incorporate up to 40 percent solvent-extracted soybean meal into commercial diets. A subsequent series of experiments identified mineral supplementation as limiting higher soybean use in hybrid striped bass. Once that was evaluated, soybean use could be increased to 45 percent to 50 percent of the diet.


Most of the evaluations with fish and crustaceans have not involved quantified EAA requirements of the targeted species. Experimental formulations are most often on a crude protein basis, a method that may be yielding suspicious results. The striped bass, a hybrid, is a strict carnivore. Thus, following the model created by evaluations

with salmonids, one might assume that hybrids would not handle high levels of soybean meal.

This has not been the case. The evaluations with hybrids were one of the few in which soybean meal was substituted as a source of EAA, rather than crude protein. Thus, the crux of the problem seems to be a basic lack of understanding of the EAA requirements of the target species, and scientists who substitute soy products simply on a crude protein basis.

Additionally, the lack of specificity in experiments with soy led to numerous speculations on the cause of soy's limited use in diets. Most focused on trypsin inhibitor activity; others on saponins and, more recently, allergenic proteins. However, the number of detailed studies examining any of these anti-nutritional factors is limited, which is somewhat surprising, considering there have been more than 200 evaluations of soy in fish and crustacean diets.





In general, the most common method used to date for evaluating soybeans in aquatic diets is to include them at increasing levels, and sometimes maintain isonitrogenous diets. Other times, soy is simply included at some level, and then the cause of any problems is speculated.


The research needed to expand the use of soy products in aquaculture requires detailed studies examining various soy components that are considered anti-nutritional factors. These data will guide researchers to appropriate processing methods to remove those components that are causing legitimate problems, or to biotechnological solutions that will result in a modified soybean for the aquaculture market.

Should soybean meal become widely used in aquatic diets, it will offer more than just a nutritional benefit. Diminishing use of fish meal and increasing use of soybean meal appears to improve the

effluent characteristics of aquaculture operations. This will be an increasingly important consideration in aquaculture, as it is in other animal production industries.

The last major food item we still hunt and gather from the wild are those animals that live underwater. However, we cannot harvest more from the seas. If man wants to consume fish and shellfish in the future, the percentage grown in commercial operations must increase. Additional research is required in order to participate in this new global industry.

The above is a synopsis of the aquaculture paper published by Paul B. Brown. The paper in its entirety, together with a list of references, can be found at www.soymeal.org.



The United Soybean Board's Better Bean Initiative and Aquaculture

by Nick Bajjalieh, Ph.D.

The United Soybean Board's Better Bean Initiative (BBI) seeks to grow demand for U.S.-produced "commodity" soybeans by improving their inherent compositional value. While the meal component of BBI has focused on swine and poultry applications, targeted improvements also offer benefit to aquaculture. BBI meal targets address the following areas:

- n Protein quality, as represented by relative amino acid levels, quantity and digestibility.
- n Carbohydrate characteristics and related nutritional energy.
- n Improved phosphorus utilization through reduced phytic acid levels.

Nutritional Considerations

PROTEIN: As with terrestrial non-ruminant animals, protein quality, quantity and digestibility are critical considerations in the formulation of diets for aquatic animals. As pointed out in Dr. Brown's article (excerpted in this newsletter), lysine and methionine are the two most limiting essential amino acids in feed formulations for most aquatic species. BBI meal trait targets include increases in both. Other protein traits targeted by BBI also represent potential benefit.

PHYTATE: The digestibility of phytate-bound phosphorus is negligible for fish. Therefore, improving the digestibility of phytate phosphorus will have nutritional and environmental benefits.



CARBOHYDRATE: The nutritional value of different types of carbohydrates varies across aquatic species. BBI changes must



be evaluated in this context. Implicit in any attempt to improve the carbohydrate characteristics of soybeans is the opportunity to gain greater knowledge about this critical, but relatively neglected component. Knowledge gained should also be of value to aquatic animal nutritionists.

Environmental Considerations

Water quality and usage is already a critical issue. It will increasingly become more so as global population continues to expand. Water quality is an especially sensitive issue where intensive aquaculture practices are employed.

Since most aquatic animals defecate directly into the aqueous environment in which they live, separation of the animals from their waste becomes a major challenge. Where abundant supplies of "clean" water are available, the concentration of waste materials can be maintained at low levels through dilution. However, this approach has both social and practical application limits as aquatic animal population densities increase.


To address this issue, various technologies such as bio-filters, creation of synergistic eco-systems, etc., continue to be developed and

refined. However, aquatic animal waste will be a major factor in the continued growth and evolution of the aquaculture industry.

While some animal waste is inevitable, the greater the extent to which a feed can be digested and productively utilized by the animal, the less its environmental impact. BBI meal trait targets focus on improving the extent and efficiency with which soybean meal serves as a source of nutrients. As such, they will help to lessen the environmental footprint associated with aquaculture's usage of soybean meal.

A Work in Progress

BBI is an ambitious, multi-faceted and ever evolving process. We are only at the early stages. For BBI to be successful, the various value chain participants must recognize the opportunity that it represents and be willing to invest as well. Through BBI, the United Soybean Board looks forward to adding value to the domestic soybean franchise.



Summary Comments

Should soybean meal become widely used in aquatic diets, it will offer more than just a nutritional benefit. Diminishing use of fish meal and increasing use of soybean meal appears to improve the effluent characteristics of aquaculture operations. This will be an increasingly important consideration in aquaculture, as it is in other animal production industries.

The United Soybean Board's Better Bean Initiative (BBI) seeks to grow demand for U.S.-produced "commodity" soybeans by improving their inherent compositional value. While the meal component of BBI has focused on swine and poultry applications, targeted improvements also offer benefit to aquaculture.



References

Paul B. Brown. March, 2002. The United Soybean Board's Better Bean Initiative and Soybean Meal in Aquaculture Industry. Soybean Meal Infosource.

N. L. Bajjalieh. March, 2002. The United Soybean Board's Better Bean Initiative and the Aquaculture Industry. Soybean Meal Infosource.

For more information, please contact:

Dr. Keith Smith
Keith Smith and Associates
15 Winchester Road
Farmington, MO 63640



The SOYBEAN MEAL INFORMATION CENTER is sponsored by the Soybean Checkoff: Including the checkoff boards from Iowa, Kansas, Michigan, Minnesota, Nebraska, North Dakota, Ohio and the United Soybean Board